

Fluorescein-Labeled Antibody To Human IgG (γ), F(ab')₂

Produced in Goat

Catalog No. Size

202-1002 1.0 mg



DESCRIPTION

F(ab')₂ fragment, purified by size exclusion from a pepsin digest of affinity purified antibody isolated from a pool of serum from goats immunized with purified human IgG was labeled with fluorescein isothiocyanate.

FORM/STORAGE

Lyophilized. Store at 2-8°C until rehydrated. Stable for a minimum of 1 year when stored at 2-8°C.

STABILIZER AND PRESERVATIVE

Goat serum and/or bovine serum albumin (BSA) is added as a protein stabilizer. No preservative added. Additional biological protection may be provided with 0.1% sodium azide. Non-sterile.

ANTIBODY CONCENTRATION

The concentration of affinity purified antibody is 1.0 mg as determined by UV absorbance at 280 nm.

F/P RATIO

The fluorescein:antibody protein ratio is between 3:1 and 7:1.

SPECIFICITY/CROSS REACTIVITY

Tested by gel diffusion and ELISA techniques as applicable. This product reacts specifically with human IgG. This antibody will not bind to the Fc receptors present on certain cell surfaces. Reactivity to IgG subclasses has not been tested. Antibodies to human IgG may cross-react with immunoglobulins of other mammalian species if common binding sites are shared. Product has been adsorbed with purified human IgA and IgM. Reactivity has not been characterized to other human immunoglobulins.

REHYDRATION AND STORAGE

Note: Rehydration of antibodies in TBS or buffers other than those listed here is not recommended.

Rehydration: Rehydrate with 1 mL of reagent quality water. Rotate the vial until the lyophilized pellet is totally dissolved.

Storage: This product may be stored for up to 1 week refrigerated; thereafter, it should be stored frozen. Stable for a minimum of 1 year at -20°C.

SUGGESTED WORKING DILUTIONS

Optimal working concentrations should be determined experimentally. Prepare working dilutions in TBS or other buffer such as 5X Detector Block or Milk/Diluent Blocking Solution (See RELATED PRODUCTS) immediately before use.

These buffers are not recommended for long term storage. A suggested starting dilution of **1:20-1:200** is recommended for most applications. In many cases, the antibody may be diluted further than indicated.

PRODUCT SAFETY AND HANDLING

This product is considered non-hazardous as defined by The Hazard Communication Standard (29 CFR 1910.1200). Avoid contact with skin and eyes. In case of contact or spillage, clean with copious amounts of water. Disposal via sanitary sewer.

RELATED PRODUCTS

Milk Diluent/Blocking Solution	Cat. No. 50-82-01
5X Detector Block	Cat. No. 71-83-00
Coating Solution Concentrate	Cat. No. 50-84-00
Wash Solution Concentrate	Cat. No. 50-63-00
10% BSA Diluent/Block	Cat. No. 50-61-00

The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Nothing disclosed herein is to be construed as a recommendation to use this product in violation of any patents. The information presented above is believed to be accurate. However, said information and product are offered without warranty or guarantee since the ultimate conditions of use and the variability of the materials treated are beyond our control. We cannot be responsible for patent infringements or other violations that may occur with the use of this product. No claims beyond replacement of unacceptable material or refund of purchase price shall be allowed. All claims regarding product performance must be made within 30 days following date of delivery.

Limited Use License

The purchase of this product conveys to the buyer the non-transferable right to use the product in research conducted by the buyer. The buyer cannot sell or otherwise transfer this product or materials made by use of this product to a third party or otherwise use this product or materials made with this product for Commercial Purposes without written approval of KPL, Inc. For additional information, please visit www.kpl.com or the KPL catalog. To obtain a license or approval to use this product for purposes other than those permitted above, contact Director of Business Development at (301) 948-7755.